



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

which give the ratios of  $x$ ,  $y$ , and  $z$  for the 8 cases corresponding to the set of internal bisectors. If we take the positive square roots in (4) and (5), we find, using the plus signs,

$$(6) \quad x : y : z = (1 + u)^{-2} : (1 + v)^{-2} : (1 + w)^{-2}$$

and, using the minus signs,

$$(7) \quad x : y : z = u^2(1 + u)^{-2} : v^2(1 + v)^{-2} : w^2(1 + w)^{-2}$$

where

$$u = \tan A/4, \quad v = \tan B/4, \quad w = \tan C/4;$$

and hence

$$(8) \quad 1 - \Sigma u - \Sigma uv + uvw = 0.$$

By (1), the actual values of  $x$ ,  $y$ , and  $z$  are  $\frac{(1+v)(1+w)}{2(1+u)}$ , etc., in the case of (6) and  $\frac{u(1+v)(1+w)}{2vw(1+u)}$ ,

etc., in the case of (7). From these two solutions we can get, not only the 8, but all the 32, by replacing  $A$ ,  $B$ ,  $C$  by  $A + l\pi$ ,  $B + m\pi$ ,  $C + n\pi$ , where  $l + m + n$  is a multiple of 4, since these angles apply equally well to the triangle, and they leave the relation (8) unaltered.

To get a practical construction, let us denote by  $\rho_1, \rho_{11}, \rho_{12}, \rho_{13}; \rho_2, \rho_{21}, \rho_{22}, \rho_{23}, \dots$ , the in- and ex-radii of  $AFI, BDI, CEI$ . Then

$$\rho_{11} = \frac{1+u}{2}r, \quad \rho_{12} = \frac{1+u}{2u}r, \text{ etc.}$$


Hence in the case of (6), the radii of the required circles are the three fourth proportionals to  $\rho_{11}, \rho_{21}, \rho_{31}$  in different orders; and in the case of (7) the fourth proportionals to  $\rho_{12}, \rho_{22}, \rho_{32}$ . The remainder of the set of 8 cases may be solved by using other combinations of the  $\rho$ 's, while all the 32 cases may be similarly treated by means of the triangles  $AF_1I_1$ , etc.

#### 495. Proposed by N. P. PANDYA, Sojitra, India.

A point  $P$  moves so that the quadrilateral  $PBCD$  is half of a given quadrilateral  $ABCD$ . Find the locus of  $P$ .

SOLUTION BY J. W. BALDWIN, University of Michigan.

In general the triangles  $ABD$  and  $BCD$  will not be equal. Let  $ABD$  be the larger of the two. Then we are to have triangle  $BCD +$  triangle  $PBD$  equal to half of the given quadrilateral  $ABCD$  for all positions of  $P$ . That is, the triangle  $PBD$  must have a constant area; and having a fixed base  $BD$  must have a constant altitude, the distance from  $P$  to  $BD$  or  $BD$  produced. Hence the locus of  $P$  is a line parallel to  $BD$ . In case triangle  $ABD =$  triangle  $BCD$  the locus of  $P$  is the line of which  $BD$  is a segment and two sides  $PB$  and  $PD$  of the quadrilateral  $PBCD$  fall in this line.



Also solved by W. J. THOME, G. W. HARTWELL, WILLIAM HOOVER, S. W. REAVES, W. R. RANSOM, J. W. CLAWSON, and the PROPOSER, some solvers using analytic methods and one using trilinear, perpendicular coördinates.

#### CALCULUS.

##### 407. Proposed by PAUL CAPRON, Annapolis, Maryland.

A coffee pot in the form of a conical frustum, 10 inches high, with a lower base 8 inches in diameter and an upper base 6 inches in diameter, is held on a slant so that the lower base is barely covered by the coffee within, and the upper base is barely uncovered. How much coffee does the pot contain?

#### III. SOLUTION BY WILLIAM W. JOHNSON, Cleveland, Ohio.

The quantity of coffee in the pot is equal to the volume of the conical ungula  $C-APBQ$  formed by tipping the conical frustum on a slant, according to the conditions of the problem. In the figure let  $AL = R$ ,  $DF = r$ ,  $FL = h$ ,  $GO = x$ ,  $FO = y$ . Then,

$$\text{Volume } C-APBQ = V = \int_r^R S dy = \frac{h}{R-r} \int_r^R \left( x^2 \arccos \frac{2Rr - (R+r)x}{(R-r)x} - \frac{1}{(R-r)^2} [2Rr - (R+r)x] \sqrt{4Rr(R+r)x - 4Rrx^2 - 4R^2r^2} \right) dx,$$

in which, area  $HKE = S$ , and

$$dy = \left( \frac{h}{R-r} \right) dx;$$

since


$$y = \frac{h(x-r)}{R-r}.$$

Integrating, we obtain

$$V = \frac{\pi h R^{\frac{3}{2}}}{3(R-r)} (R^{\frac{3}{2}} - r^{\frac{3}{2}}).^1$$

Putting  $h = 10$ ,  $R = 4$ , and  $r = 3$ , we get

$$V = \frac{80}{3} (8 - 3\sqrt{3})\pi = 234.895 \text{ cu. in.}$$



#### 409. Proposed by B. J. BROWN, Victor, Colorado.

Integrate the equation

$$\frac{\partial^2 z}{\partial x \partial y} + \frac{1}{x+y} \left( \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} \right) - \frac{2}{(x+y)^2} z = 0.$$

SOLUTION BY O. S. ADAMS, Coast and Geodetic Survey, Washington, D. C.

Let

$$(1) \quad z = \frac{u}{(x+y)^2}.$$

Then, computing  $\partial z/\partial x$ ,  $\partial z/\partial y$  and  $\partial^2 z/\partial x \partial y$ , and substituting in the given equation, we may write the result as follows,

$$\frac{\partial}{\partial y} \left( \frac{\partial u}{\partial x} - \frac{u}{x+y} \right) - \frac{1}{x+y} \left( \frac{\partial u}{\partial x} - \frac{u}{x+y} \right) - \frac{2u}{(x+y)^2} = 0.$$

Now let

$$(2) \quad \frac{\partial u}{\partial x} - \frac{u}{x+y} = v.$$

Then

$$(3) \quad u = \frac{(x+y)^2}{2} \frac{\partial v}{\partial y} - \frac{x+y}{2} v.$$

Computing  $\partial u/\partial x$  and  $u/(x+y)$ , and substituting in (2), we have

$$\frac{\partial u}{\partial x} - \frac{u}{x+y} = \frac{(x+y)^2}{2} \frac{\partial^2 v}{\partial x \partial y} + \frac{x+y}{2} \frac{\partial v}{\partial y} - \frac{x+y}{2} \frac{\partial v}{\partial x} = v;$$

or

$$\frac{\partial^2 v}{\partial x \partial y} + \frac{1}{x+y} \frac{\partial v}{\partial y} - \frac{1}{x+y} \frac{\partial v}{\partial x} - \frac{2v}{(x+y)^2} = 0.$$

This equation may be written in the form,

$$\frac{\partial}{\partial y} \left( \frac{\partial v}{\partial x} + \frac{v}{x+y} \right) - \frac{1}{x+y} \left( \frac{\partial v}{\partial x} + \frac{v}{x+y} \right) = 0.$$

Finally, let

$$(4) \quad \frac{\partial v}{\partial x} + \frac{v}{x+y} = w.$$

Then

$$\frac{\partial w}{\partial y} - \frac{w}{x+y} = 0; \quad \text{or} \quad \frac{1}{x+y} \frac{\partial w}{\partial y} - \frac{w}{(x+y)^2} = 0,$$

<sup>1</sup> See Finkel's *Solution Book*, p. 319.